

RP00992

Leader in Biomolecular Solutions for Life Science



Recombinant Human TNFRSF10D/DcR2/TRAIL-R4/CD264 Protein

Catalog No.: RP00992

Recombinant

Sequence Information

Species	Gene ID	Swiss Prot
Human	8793	Q9UBN6

Tags

C-hFc&His

Synonyms

CD264; DCR2; TRAIL-R4; TRAILR4; TRUNDD; TNFRSF10D; DCR2; TRAIL-R4; TRAILR4; TRUNDD

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE.

Endotoxin

< 0.1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize freeze-thaw cycles.

Background

Basic Information

Description

Active Recombinant Human TNFRSF10D/DcR2/TRAIL-R4 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ala 56 - His 211) of human DcR2 (Accession #NP_003831.2) fused with an Fc, 6×His tag at the C-terminus.

Bio-Activity

1. Measured by its binding ability in a functional ELISA. Immobilized Recombinant Human TRAIL at 2 μg/mL (100 μL/well) can bind Recombinant Human DcR2 with a linear range of 15-60 ng/mL. 2. Measured by its ability to inhibit TRAIL-mediated cytotoxicity using L-929 mouse fibroblast cells treated with TRAIL. The ED₅₀ for this effect is 1.8-7.2 ng/mL in the presence of 20 ng/mL Recombinant Human TRAIL/TNFSF10.

Storage


Store the lyophilized protein at -20°C to -80 °C for long term. After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week. Avoid repeated freeze/thaw cycles.

Contact




www.abclonal.com


Validation Data



Active Recombinant Human TNFRSF10D/DcR2/TRAIL-R4 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 60-70 kDa.



Immobilized Recombinant Human TRAIL at 2 μ g/mL (100 μ L/well) can bind Recombinant Human DcR2 with a linear range of 15-60 ng/mL.



Recombinant Human TNFRSF10D inhibit TRAIL-mediated cytotoxicity using L-929 mouse fibroblast cells treated with TRAIL. The ED₅₀ for this effect is 1.8-7.2 ng/mL in the presence of 20 ng/mL Recombinant Human TRAIL/TNFSF10.