

RP01064

Leader in Biomolecular Solutions for Life Science



# Recombinant Mouse Lipocalin-2/NGAL/LCN2 Protein

Catalog No.: RP01064

Recombinant

## Sequence Information

Species	Gene ID	Swiss Prot
Mouse	16819	P11672

### Tags

C-His

### Synonyms

NGAL;LCN2;Lipocalin-2;24p3;MSFI;LCN2

## Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE.

### Endotoxin

< 0.1 EU/μg of the protein by LAL method.

### Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize freeze-thaw cycles.

## Contact



[www.abclonal.com](http://www.abclonal.com)

## Background

### Basic Information

#### Description

Recombinant Mouse Lipocalin-2/NGAL Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Gln21-Asn200) of mouse Lipocalin-2/NGAL (Accession #NP\_032517.1) fused with an 8×His tag at the C-terminus.


#### Bio-Activity

Measured by its ability to bind Iron(III) dihydroxybenzoic acid [Fe(DHBA)3]. The binding of Fe(DHBA)3 results in the quenching of Trp fluorescence in recombinant mouse Lipocalin-2. Recombinant mouse Lipocalin-2 can bind >47.7 μM of Fe(DHBA)3.


#### Storage

Store the lyophilized protein at -20°C to -80 °C for long term. After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week. Avoid repeated freeze/thaw cycles.

## Validation Data



Recombinant Mouse Lipocalin-2/NGAL/LCN2 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 26-27kDa.



The binding of Fe(DHBA)3 results in the quenching of Trp fluorescence in recombinant mouse Lipocalin-2. Recombinant mouse Lipocalin-2 can bind >47.7 μM of Fe(DHBA)3.