Leader in Biomolecular Solutions for Life Science

ABclonal® www.abclonal.com

Recombinant Mouse Latent TGF-beta 1 Protein

Catalog No.: RP01167 Recombinant

Sequence Information

Species Gene ID Swiss Prot Mouse 21803 P04202

Tags

N-His

Synonyms

TGF-

beta1;Tgfb;Tgfb-1;TGFbeta1;CED;DPD1;L AP;TGFB;TGFbeta;TGFB1

Product Information

Source Purification HEK293 cells > 95% by SDS-PAGE.

Endotoxin

 $< 0.1 \; \text{EU/}\mu\text{g}$ of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile 4 mM HCl. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact



www.abclonal.com

Background

Basic Information

Description

Recombinant Mouse Latent TGF-beta 1 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Leu30-Ser390) of mouse TGF-beta 1 (Accession #NP 035707.1) fused with a 8×His tag at the N-terminus.


Bio-Activity

Measured by its ability to inhibit the IL-4-dependent proliferation of HT-2 mouse T cells. The ED₅₀ for this effect is 0.02-0.1 ng/mL, corresponding to a specific activity of 1.0×10 ⁷- 5.0×10 ⁷-units/mg.


Storage

Store the lyophilized protein at -20°C to -80 °C for long term.
br>After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week. Avoid repeated freeze/thaw cycles.

Validation Data



Active Recombinant Mouse TGF-beta 1 Protein was determined by SDS-PAGE with Coomassie Blue, showing bands at 14 kDa, 40-50 kDa and 55 kDa..



Recombinant Mouse TGF-beta 1 inhibits the IL-4-dependent proliferation of HT-2 mouse T cells. The ED $_{50}$ for this effect is 0.02-0.1 ng/mL, corresponding to a specific activity of 1.0×10^7 - 5.0×10^7 units/mg.