Leader in Biomolecular Solutions for Life Science

ABclonal www.abclonal.com

Recombinant Mouse Fc-gamma RIV/CD16-2 Protein

Catalog No.: RP01394 Recombinant

Sequence Information

Species Gene ID Swiss ProtMouse 246256 A0A0B4|1G0

Tags

C-His

Synonyms

Fc gamma RIV;CD16-2;Fcgr4;FCGR4;mouse igg;FcgR4

Product Information

Source

Purification

HEK293 cells

> 95% by SDS-PAGE.

Endotoxin

<0.1EU/µg

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact



www.abclonal.com

Background

FcgammaRIV is a relatively new IgG Fc receptor (FcgammaR) that is reported to contribute to the pathogenesis of autoimmune diseases.FcgammaRIII and FcgammaRIV are each essential to trigger an FcRgamma-linker for activation of T-cell-dependent signal that drives C5a production in the Arthus reaction. A combined requirement for FcgammaRIII and FcgammaRIV in autoimmune injury, and identify the linker for activation of T cells adaptor as an integral component of linked FcgammaR and C5a anaphylatoxin receptor activation to generate inflammation.

Basic Information

Description

Recombinant Mouse Fc-gamma RIV/CD16-2 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gly21-Gln 203) of mouse Fc gamma RIV/CD16-2 (Accession $\#NP_653142.2$) fused with a 6×His tag at the C-terminus.

Bio-Activity


Storage

Store the lyophilized protein at -20°C to -80°C for long term.

-After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Mouse Fc-gamma RIV/CD16-2 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 32-38kDa.