

RP01443

Leader in Biomolecular Solutions for Life Science



Recombinant Human Apolipoprotein E/ApoE Protein

Catalog No.: RP01443

Recombinant

Sequence Information

Species	Gene ID	Swiss Prot
Human	348	P02649

Tags

C-His

Synonyms

APOE;AD2;APO-E;ApoE4;LDLCQ5;LPG;ApoE

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE.

Endotoxin

<0.1EU/μg

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

This protein is a major apoprotein of the chylomicron. It binds to a specific liver and peripheral cell receptor, and is essential for the normal catabolism of triglyceride-rich lipoprotein constituents. This gene maps to chromosome 19 in a cluster with the related apolipoprotein C1 and C2 genes. Mutations in this gene result in familial dysbetalipoproteinemia, or type III hyperlipoproteinemia (HLP III), in which increased plasma cholesterol and triglycerides are the consequence of impaired clearance of chylomicron and VLDL remnants.

Basic Information

Description

Recombinant Human Apolipoprotein E/ApoE Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Lys19-His317) of human Apolipoprotein E/ApoE (Accession #NP_000032.1) fused with a 6×His tag at the C-terminus.

Bio-Activity

Storage


Store the lyophilized protein at -20°C to -80°C for long term. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.

Contact




www.abclonal.com

Validation Data



Recombinant Human Apolipoprotein E/ApoE Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 36-38kDa.



Immobilized Human ApoE at 5μg/mL (100 μL/well) can bind Human TREM2 with a linear range of 0.6-65 ng/mL .