Leader in Biomolecular Solutions for Life Science



Biotinylated Recombinant Human VEGF-A/VEGF165 Protein

Catalog No.: RP02097 Recombinant 1 Publications

Sequence Information

Species Gene ID Swiss Prot Human 7422 P15692-4

Tags

C-His&Avi

Synonyms

VEGFA; MVCD1; VEGF; VPF; vascular endothelial growth factor A;MVCD1;VEGF;VPF;L VEGFA;VEGF A

Product Information

Source Purification
HEK293 cells > 95% by Tris-Bis
PAGE

Endotoxin

< 1 EU/µg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact



www.abclonal.com

Background

Basic Information

Description

Biotinylated Recombinant Human VEGF165 Protein is produced by Expi293 expression system. The target protein is expressed with sequence (Ala27-Arg191) of Human VEGF165 fused with a His tag and Avi tag at the C-terminal.

Bio-Activity

Immobilized biotinylated Human VEGFR1 at 0.5 μ g/mL (100 μ L/Well).Dose response curve for Biotinylated Human VEGF165 with the EC₅₀ of 26.17 ng/mL determined by ELISA.


Storage

Store the lyophilized protein at -20°C to -80°C for long term.


-After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Validation Data



Biotinylated VEGF165 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of Biotinylated Human VEGF165 is greater than 95% as determined by SEC-HPLC.



Immobilized Biotinylated Human VEGF165 at $0.2\mu g/ml$ (100 $\mu l/Well$) on the plate. Dose response curve for Anti-VEGF165 Antibody, hFc Tag with the EC₅₀ of 11.3ng/ml determined by ELISA.