

RP02324

Leader in Biomolecular Solutions for Life Science



Recombinant SARS-COV-2 Spike S1(N501Y) Protein

Catalog No.: RP02324

Recombinant

Sequence Information

Species	Gene ID	Swiss Prot
SARS-COV-2		YP_00972439
		0.1

Tags

C-His

Synonyms

S1 protein;Spike glycoprotein Subunit1;S glycoprotein Subunit1;Spike protein S1

Product Information

Source	Purification
HEK293 cells	> 95% by Tris-Bis PAGE;> 95% by SEC-HPLC

Endotoxin

< 1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

Basic Information

Description

Recombinant SARS-COV-2 SARS-COV-2 Spike S1 (N501Y) Protein is produced by Expi293 expression system. The target protein is expressed with sequence (Val16-Arg685) of SARS-COV-2 SARS-COV-2 Spike S1 (N501Y) fused with His tag at the C-terminal.

Bio-Activity

Immobilized SARS-COV-2 Spike S1 (N501Y) ,His Tag at 0.5 μg/mL (100 μL/Well) on the plate. Dose response curve for Human ACE2 ,hFc Tag with the EC₅₀ of 80.3 ng/mL determined by ELISA.

Storage


Store the lyophilized protein at -20°C to -80°C for long term.
After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.

Contact




www.abclonal.com


Validation Data



SARS-COV-2 Spike S1 (N501Y) on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of SARS-COV-2 Spike S1 (N501Y) is greater than 95% as determined by SEC-HPLC.



Immobilized SARS-COV-2 Spike S1 (N501Y) , His Tag at 0.5μg/ml (100μl/Well) on the plate. Dose response curve for Human ACE2 , hFc Tag with the EC₅₀ of 80.3ng/ml determined by ELISA.