Leader in Biomolecular Solutions for Life Science



Biotinylated Recombinant Human Siglec-10 Protein

Catalog No.: RP02499 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 89790 Q96LC7

Tags

C-hFc&Avi

Synonyms

SLG2; PRO940; SIGLEC-10; SIGLEC10

Product Information

Source HEK293 cells

Purification > 95% by Tris-Bis

PAGE;> 95% by SEC-HPLC

Endotoxin

< 1 EU/µg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact



www.abclonal.com

Background

Basic Information

Description

Biotinylated Recombinant Human siglec 10 Protein is produced by Expi293 expression system. The target protein is expressed with sequence (Met17-Thr546) of Human siglec 10 fused with hFc and Avi tag at the C-terminal.


Bio-Activity

Storage


Store the lyophilized protein at -20°C to -80°C for long term.

h/>After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.


Validation Data



Biotinylated Human Siglec-10 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of Biotinylated Human Siglec-10 is greater than 95% as determined by SEC-HPLC.



Immobilized Anti-Siglec-10 Antibody, hFc Tag at 1μ g/ml (100μ l/well) on the plate. Dose response curve for Biotinylated Human Siglec-10, hFc Tag with the EC₅₀ of 25ng/ml determined by ELISA.