

RP02547

Leader in Biomolecular Solutions for Life Science



Biotinylated Recombinant Human FGFR3 beta (IIIc)/CD333 Protein

Catalog No.: RP02547

Recombinant

Sequence Information

Species	Gene ID	Swiss Prot
Human		P22607

Tags

C-His&Avi

Synonyms

ACH; CD333; CEK; CEK2; EC 2.7.10; FGF R3; FGFR3; HSFGR3EX; JTK4

Background

Four distinct genes encoding closely related FGF receptors, FGF R1-4, are known. All four genes for FGF Rs encode proteins with an N-terminal signal peptide, three immunoglobulin (Ig)-like domains, an acid?box region containing a run of acidic residues between the Igl and IgII domains, a transmembrane domain and the split tyrosine-kinase domain. FGFR3 is tyrosine-protein kinase that acts as cell-surface receptor for fibroblast growth factors and plays an essential role in the regulation of cell proliferation, differentiation and apoptosis. Plays an essential role in the regulation of chondrocyte differentiation, proliferation and apoptosis, and is required for normal skeleton development. Regulates both osteogenesis and postnatal bone mineralization by osteoblasts..

Product Information

Source

HEK293 cells

Purification

> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Endotoxin

Less than 1EU per µg by the LAL method.

Formulation

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Basic Information

Description

Recombinant Biotinylated Human FGFR3 beta (IIIc) Protein is expressed from Expi293 with His tag and Avi tag at the C-terminal. It contains Asp127-Gly377.

Bio-Activity

Storage


Store the lyophilized protein at -20°C to -80°C for long term. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.

Contact




www.abclonal.com


Validation Data



Biotinylated Human FGFR3 beta (IIIC) on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of Biotinylated Human FGFR3 beta (IIIC) is greater than 95% as determined by SEC-HPLC.



Immobilized Biotinylated Human FGFR3 beta (IIIC) at 0.5µg/ml (100µl/Well) on the plate. Dose response curve for Anti-FGFR3 Antibody, hFc Tag with the EC₅₀ of 11.4ng/ml determined by ELISA.