RP02584

Leader in Biomolecular Solutions for Life Science

# **Biotinylated Recombinant Human IL-12A&IL-12B Protein**



Catalog No.: RP02584 Recom

4 Recombinant

## Sequence Information

 Species
 Gene ID
 Swiss Prot

 Human
 3592&3593
 P29460(IL-12

 B)&P29459(I
 L-12A)

Tags

C-His&Avi

## Synonyms

CLMF; CLMF2; IL-12A; IL-12B; IL12; IL12 p70; IMD28; IMD29; NFSK; NKSF1; NKSF2; P35; IL-12 subunit p35; IL12A; interleukin 12

## **Product Information**

Source	Purification
HEK293 cells	> 95% as
	determined by Tris-
	Bis PAGE□> 95% as
	determined by HPLC

## Endotoxin

Less than 1EU per  $\mu$ g by the LAL method.

## Formulation

## Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## Contact

www.abclonal.com

Background

Interleukin 12 (IL-12) is a pleiotropic cytokine that plays an essential role in Th1-type immune response against cancer, a condition where cells in a particular part of the body grow and reproduce uncontrollably.

# **Basic Information**

## Description


Recombinant Biotinylated Human IL-12 Protein is expressed from Expi293 with His tag and Avi tag at the C-terminal. [It contains Ile23-Ser328(IL-12B)/Arg23-Ser219(IL-12A).

## **Bio-Activity**


## Storage

Store the lyophilized protein at -20°C to -80°C for long term.<br/>After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.

# Validation Data



The purity of Bitinylated Human IL-12 is greater than 95% as determined by SEC-HPLC.



Immobilized Anti-IL-12 Antibody, hFc Tag at 0.5 $\mu$ g/ml (100 $\mu$ I/well) on the plate. Dose response curve for Biotinylated Human IL-12, His Tag with the EC<sub>so</sub> of 9.3ng/ml determined by ELISA.

Biotinylated Human IL-12 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.